

## DESCRIPTION

### MATERIALS AND PROCEDURES FOR THE PURIFICATION OF CELLS

5

#### Cross-Reference to Related Applications

This application is a continuation of co-pending application U.S. Serial No. 09/437,509, filed November 10, 1999, which is a continuation of application U.S. Serial No. 08/965,949, filed November 7, 1997, now abandoned, which claims the benefit of provisional application U.S. Serial No. 60/030,725, filed November 8, 1996.

10

#### Background of the Invention

Mammalian blood cells provide for an extraordinarily diverse range of activities. Hematopoietic stem cells are defined as those cells that are capable of both self-renewal and differentiation into the two principle precursor components -- the myeloid and lymphoid lines. Such stem cells are said to be "totipotent." Stem cells that are less general but that can still differentiate into several lines are called "pluripotent." Further differentiation then occurs among the precursor cells to produce the monocyte, eosinophil, neutrophil, basophil, megakaryocytes, and erythroid lineages from the myeloid line, and T cells, B cells, and NK cells from the lymphoid line. Hematopoietic stem cells reside primarily in the bone marrow.

15

20

25

One of the first breakthroughs into stem cell isolation and identification came in the late 1980's. In U.S. Patent No. 4,714,680, Civin described pluripotent lympho-hematopoietic cells that were substantially free of mature lymphoid and myeloid cells. Civin also described an antigen, MY-10, and a monoclonal antibody thereto, which was present on those cells. Those cells made up about 1% of all cells in normal adult bone marrow, and generally comprised a mixture of totipotent, and pluripotent stem cells and lineage committed precursor cells with the latter cells predominating. Since that time, MY-10 has been classified by the International Workshop on Human Leukocyte Antigens as falling with the cluster designated

as "CD34." Anti-CD34 monoclonal antibodies are now commercially available from a number of sources including, for example, Becton Dickinson Immunocytometry Systems ("BDIS").

Other investigators have attempted to subset CD34<sup>+</sup> cells from both peripheral blood and bone marrow. Bender *et al.*, *Blood* 77:2591-2596 (June 1991), used four color flow cytometry with combinations of monoclonal antibodies (*i.e.*, anti-CD34, anti-CD33, anti-CD45, anti-CD19, anti-CD7, anti-CD10, anti-CD3, anti-CD20, anti-CD14, anti-CD11b and anti-HLA-DR), to identify and isolate CD34<sup>+</sup> hematopoietic progenitor cells.

There is a strong interest in identifying and isolating human hematopoietic stem cells. However, the stem cell population constitutes only a small percentage of the total number of leukocytes in bone marrow. In view of the small proportion of the total number of cells in the bone marrow which are stem cells, the uncertainty of the markers associated with the stem cell as distinct from more differentiated cells, and the general inability to biologically assay for human stem cells, the identification and purification of stem cells has been difficult.

Having a procedure for the efficient isolation of stem cells would allow for identification of growth factors associated with, for example, (1) the early steps of dedication of the stem cell to a particular lineage; (2) the prevention of such dedication; and (3) the negative control of stem cell proliferation. Readily available stem cells would also be extremely useful in bone marrow transplantation, as well as transplantation of other organs in association with the transplantation of bone marrow. Also, stem cells are important targets for gene therapy, where the inserted genes promote the health of the individual into whom the stem cells are transplanted. In addition, the ability to isolate stem cells may serve in the treatment of lymphomas and leukemia, as well as other neoplastic conditions. The identification and isolation of the most primitive population of hematopoietic stem cells would be highly advantageous in situations where reinfusion of only a small number of long-term repopulating cells was desired. For example, this would be the case when purging bone

marrow or peripheral blood stem cells of contaminating tumor cells, or where genetic manipulation of the stem cells was the objective.

The separation of a particular mammalian cell population from a mixture of cell populations is quite different from the separation of chemical species such as proteins from a solution. Most mammalian cells are on the order of 8 to 20 microns ( $\mu$ ) in diameter. In contrast, the proteins and other chemical species are significantly smaller, *i.e.*, on the order of 1000-fold or more. In addition, another factor unique to the separation of mammalian cells is the need to preserve cell viability.

In contrast to yeast cells, which are relatively insensitive to changes in osmolarity, pH and shear, higher order mammalian cells are much more sensitive to shear forces-exerted during purification, pH osmolarity, and the chemical composition of the reagents used. Therefore, the steps comprising the method and all reagents used must be non-toxic to the cells.

Separation of mixtures of chemicals, biomolecules and cell types is often effected by immunoaffinity chromatography. Packed beds, such as those used in column chromatography, are often used in affinity separation. However, problems such as non-specific trapping or filtration of cells and clogging make the use of a packed bed undesirable for cell separation.

One device that has been developed for reducing the pressure drop across a column of particles is the fluidized bed. A fluidized bed consists of solid particles and a gas or liquid which is passed upwardly through the particle bed with velocity sufficient to fluidize the bed. The fluidization of the bed provides more surface contact between the particle and the fluid passing through the bed. One disadvantage associated with fluidized beds is the radial and axial movement of the particles which result in significant intermixing of the particles.

A number of other methods have been developed for fractionating heterogeneous mixtures of cells into the various compartments. These methods are based on the size and density of the cells, specific binding properties that they possess, and their expression of

surface antigens. The method chosen usually depends on the degree of purity required, the intended use of the selected cells, and the abundance of the cells of interest.

Density gradient centrifugation, velocity sedimentation, and counterflow centrifugal elutriation are methods currently used to separate cells based on their physical properties such as size and density. While these methods work well as pre-enrichment steps, none are accurate and/or specific enough to yield pure populations of stem cells.

Flow cytometry is extremely sensitive because it looks at each cell individually. It can distinguish multiple markers, their relative level of expression, the size and granularity of each cell, and can sort out specific cells into a waiting tube. However, the equipment is highly sophisticated, the processing is relatively slow, and it is difficult to sterilize the instrument between samples.

By immobilizing the antibody on a solid phase, several methods have been used to process larger cell numbers in a relatively short time while still exploiting the specificity of the antigen/antibody interaction. Panning is an effective three-step technique for cell selection. First, mononuclear cells are separated from bone marrow, peripheral, or cord blood by density gradient centrifugation, negatively panned for soybean agglutinin binding, and positively panned with an anti-CD34 antibody-coated flask. The selected cells are released by mechanical agitation. The major problems with this method are the multiplicity of steps, subsequent low yields and the surface area required to give clinically useful stem cell numbers.

Another method along these lines uses magnetic beads as a solid support. Mononuclear cells are incubated with anti-CD34 antibody and bound to magnetic beads coated with sheep anti-mouse antibodies. The CD34+ cells are pulled to the side of the tube by applying a magnet. Historically, it has been difficult to dislodge the beads from the cells without harming the cells and perhaps of greater concern is that incomplete removal of the beads prior to reinfusion may be harmful to the patient.

One clinical technique is cell separation by avidin-biotin column chromatography as developed by Berenson *et al.* White blood cells incubated with a biotinylated anti-CD34

antibody are applied to a column containing avidin-coated polyacrylamide or agarose beads. After washing, the bound CD34<sup>+</sup> cells are stripped from the bed by mechanical agitation. Theoretically, the mechanical release breaks the link at its weakest point, the antigen/antibody, and leaves the CD34 antigen intact on the cell surface. This is proposed because of the high affinity between biotin and avidin. Cells selected from bone marrow, peripheral blood, and cord blood by this method have been allogeneically (unrelated donor) or autologously (self) transplanted into more than 200 patients worldwide to treat breast cancer, neuroblastoma, non-Hodgkin's lymphoma, and chronic myelogenous leukemia.

There are a number of innate problems facing any rare cell separation system. For instance, large surfaces tend to bind cells nonspecifically, thus decreasing specificity. Whereas low specificity is easy to overcome in systems where large numbers of the desired cells are available, it becomes a critical factor in separating stem cells since they are in such low abundance and in a heterogeneous cell background. Detachment is also a problematic stage in separation procedures. Detachment methods currently employed include the use of enzymes that chemically clip the cells from the solid phase, mechanical methods which tear them off, and polyclonal antibodies which compete them off.

A highly purified population of stem cells is necessary for a variety of *in vitro* experiments and *in vivo* indications. For instance, a purified population of stem cells will allow for identification of growth factors associated with their self-regeneration. In addition, there may be as yet undiscovered growth factors associated (1) with the early steps of dedication of the stem cell to a particular lineage; (2) the prevention of such dedication; and (3) the negative control of stem cell proliferation.

A comparison between current cell selection methods has proven that the use of antibodies to target specific cell populations consistently results in higher specificity and recovery when compared to non-antibody based methods. However, the current methods cannot produce pure cell populations (purity  $\leq$  70%) and cannot recover more than 65% of the target cells (de Wynter *et al.*, 1995). All current methods require the sample to be

collected, often frozen and transferred to the laboratory for cell selection/depletion. None of the existing devices has the capacity to perform “in line” cell selection, that is, to directly select cells from normal peripheral blood as it is being drawn from a donor. Such possibility would make stem cell donation as simple as blood donation and would have enormous implications in the field of stem cell transplantation. It would tremendously increase the number of potential donors and expand the histocompatibility repertoire of stem cells available for transplantation, probably making the concept of stem cell banks a reality. The subject invention provides an efficient system for stem cell purification.

#### Brief Summary of the Invention

The subject invention pertains to materials and methods for target molecule and cell separation and, in particular, stem cell separation, using a polymer system. In a preferred embodiment the polymer is a conductive polymer. In one embodiment, an immunopolymer useful according to the subject invention comprises a conductive solid matrix wherein antibodies are entrapped. Preferably, these antibodies are monoclonal antibodies. The polymer with the antibodies is then contacted with a composition containing stem cells. The stem cells become associated with the antibodies within the polymer while the rest of the composition, including irrelevant cells, and other material passes through the system. The stem cells can then be released from the polymer by application of an appropriate current.

The use of an electric current to induce the release of bound cells from the immunopolymer surface provides a reliable, reproducible, and gentle method to harvest unaltered selected cells. Also, since this technique permits the monitoring of cell binding to the solid matrix during the selection process, it is possible to obtain feedback information on the number of selected cells at any time.

#### Brief Description of the Drawings

**Figure 1** shows a system according to the subject invention for purifying stem cells.

**Figure 2** shows capacity of 2 different types of conductive immunopolymers to bind CD34<sup>+</sup> cells as a function of increasing amounts of anti-CD34 antibody present during the polymerization reactions.

**Figure 3** shows effect of thickness of the membranes on cell binding capacity of the 2 types of conductive immunopolymers.

**Figure 4** shows effect of time of polymerization (+ or - anti-CD34 antibody) on the cell binding capacity of the 2 types of polypyrrole polymers.

**Figure 5** shows a reaction diagram illustrating the coupling of the enzymes HRP and GOx.

#### Detailed Disclosure of the Invention

The methods of the subject invention involve the use of molecules such as a protein, antibody or nucleic acid capable of binding to a target moiety. Exemplified herein is the use of cell-specific antibodies combined with a polymer. In accordance with the teachings of the subject invention, the antibodies are trapped inside the polymer matrix as it is formed without loss of their binding functions. A complex mixture of cells and other materials can then be passed over the polymer-antibody complex, unbound cells and other materials are removed by washing, and the cells that are trapped by the antibody can be released into a collection tube by reversing the conditions of polymer formation. Specifically, an electric current used to make a polymer can be reversed thus resulting in the release of the desired cells.

The major advantages of the system of the subject invention include the stability of the polymer-antibody complex; the polymer itself does not bind unwanted cells like the beads and panning surfaces used in other systems; and the method of detachment is fast, easy, and it does not harm the cells.

Conductive immunopolymers represent a novel approach to immunoaffinity and cell selection. In one embodiment, conductive immunopolymers are synthesized by electrically inducing polymerization of organic monomers in the presence of antibodies. This process

results in the entrapment of monoclonal antibodies into a conductive solid matrix designated as a conductive immunopolymer. None of currently available cell selection devices integrate this type of conductive solid matrix.

Conductive polymers can also be formed using enzyme generated means. In one embodiment, a polymer can be formed using an enzyme generated conducting polyphenol polymer system. Horseradish peroxidase (HRP) can be used to catalyze the polymerization of a stable electroactive phenolic conducting polymer onto the surface of a glass electrode (Alva *et al.*, 1996; the disclosure of which is herein incorporated by reference).

Incorporation of an antibody into enzyme generated polymers allows not only capture and isolation of cell populations, but also, through electrochemical measurements, provides for the monitoring of both binding and release of captured cells. Enzyme generated conductive polymers offer the following advantages as well: (1) it is a one step process; (2) there are no special purification steps; (3) the environment is totally aqueous; (4) the uncharged polymer allows for the design of positively or negatively charged, pH-type titrateable functional groups into the membrane that can assist in cell isolation; and (5) provides for the addition of other enzymes to the system such that the enzyme electrode can act as a potential specific biosensor that can be utilized in the development of monitoring systems for cell binding, release and viability.

The ability to monitor the cell binding to the solid matrix during the selection process has never before been an option. The use of a conductive immunopolymer provides this advantageous capability. The number of cells bound to the conductive polymer can be monitored by measuring the sum of the micro currents induced by each cell when maintained in immediate proximity of the polymer surface via the antibody. In the context of an "on line" separation device, this feedback information on the number of cells selected at any one time increases the efficiency and safety of the selection process since it allows the monitoring of the procedure by a computer or an operator.



The complete and gentle release of matrix bound cells represent another challenge for today's cell selection devices. Most current positive selection procedures yield altered selected cells since antibodies are still bound to their surface. The use of an electric current to induce the release of bound cell from the immunopolymer surface provides a reliable, reproducible and gentle method to harvest unaltered selected cells.

The binding capacity and specificity of the immunopolymer are primarily determined by the concentrations and specificities of the antibodies included in the polymerization mixture. The rapid and efficient selection of rare cells in large cell mixtures requires that cells interact with vast surface areas of immunopolymers. Beads represent an efficient way to present large immunoselective surface areas to a cell mixture, however, such structure is not compatible with the use of the conductive properties of the immunopolymer. Therefore, in a preferred embodiment, the subject invention utilizes polymer structures such as capillaries, spiral sheets, and parallel sheets with large surface areas and consistent conductive properties.

Non-specific binding is a critical factor limiting the specificity of the current cell selection devices. In order to electrostatically minimize non-specific binding, the immunopolymer can be designed to expose a negative net charge at the cell-matrix interface. This negative charge needs to be strong enough to prevent binding of irrelevant cells but weak enough to not interfere with the specific interaction between antibodies and target cells. Such negative charge can be a direct consequence of the polymer chemical structure or induced by a current. The latter possibility offers flexibility since it allows the operator/computer to adjust the net charge of the polymer surface to various samples and selection conditions.

Also, the prospect of a computer controlled device for the "in line" selection of stem cells from whole blood provides a very significant improvement in terms of ease of use, reliability, reproducibility and safety. Only a conductive material can allow a direct interface between electronic controls and cell selection device. This direct interface results in a more reliable, cost effective and simpler design.

The subject invention is exemplified herein using certain conductive polymers with specific antibodies. For example, antibodies to CD34 are specifically exemplified herein. However, those skilled in this art, having the benefit of the instant disclosure could readily utilize other conductive polymers and/or antibodies as described herein to effect the desired stem cell purification. For example, anti-CD4 and anti-CD8 antibodies can be used to purify sub-populations of T lymphocytes that express either CD4 or CD8 determinants.

Suitable antibodies for use in the subject invention include monoclonal and polyclonal antibody compositions. Preferably, the antibodies used in the subject invention are monoclonal antibodies. Whole antibodies and antigen binding fragments thereof are contemplated in the present invention. Thus, for example, suitable antigen binding fragments include Fab<sub>2</sub>, Fab and Fv antibody fragments.

Antibodies can also be incorporated into the conductive polymer by indirect means. For example, receptors that bind the Fc portion of an antibody molecule can be incorporated into the conductive polymer, then selected antibody can be contacted with the Fc receptor incorporated into the polymer. Antibody is then bound by the Fc receptor. Using this method advantageously provides for orienting the antibody on one particular side of the polymer matrix. Protein G, as well as other antibody binding molecules, can be used according to the subject invention.

Other indirect means can be used to attach antibody to the polymer matrix. In an exemplified embodiment, an avidin-related preparation is incorporated into the polymer matrix. Biotin-labeled antibody, either free or pre-bound to target cells, can then be contracted with and bound by the polymer through the interaction of the avidin-biotin complex formed.

Examples of suitable electrically-conducting polymers include polyheteroaromatic polymers, polyacetylenes, polyparaphenylenes, polyphenylenevinylenes and derivatives thereof. Preferred substrates include polyheteroaromatic polymers. The term "polyheteroaromatic polymer," as defined herein, means an aromatic polymer having

repeating units which include at least two elements in the aromatic rings of the repeating units.

Although the mechanism is not completely understood, it is believed that suitable polymers are electrically conductive because electrochemical oxidation, for example, causes positively charged regions, or “defects,” which allow electrons to move along polymer chains which include such defects, thereby conducting an electrical current. Examples of suitable polyheteroaromatic polymers include alkyl-substituted polythiophenes and polypyrrole. A particularly preferred polyheteroaromatic polymer is polypyrrole.

The polymer substrate is formed by, for example, electrochemical polymerization. In one example of electrochemical polymerization, a polypyrrole film is formed by disposing a graphite-rod cathode and a platinum-plate anode into a monomer solution of distilled pyrrole. An appropriate current density is maintained at a suitable voltage for a sufficient period of time to cause a polypyrrole film to form on the anode.

A further aspect of the subject invention pertains to devices useful in purifying stem cells using the procedures of the subject invention. One system for purifying stem cells is shown in Figure 1. As shown in this Figure, bone marrow can be passed through membranes of conductive polymers with which antibodies are associated. In a preferred embodiment, the antibodies would be to CD34. As shown in Figure 1, unbound material passes through the membranes and is discarded or collected for further processing. Cells which are trapped by the polymer/antibody matrix are then released by passing an appropriate current through the polymer. The cells thus released are collected for use in any one of the many applications for purified stem cells.

The subject invention can be used to simultaneously and/or sequentially collect and fractionate various cell types. This can be accomplished by utilizing antibodies with varying specificities for the different cell types. For example, the method of the subject invention can be practiced utilizing antibodies to a various cells associated with the immune system such as T cells, natural killer (NK) cells, B cells, and others. In a specific embodiment, these cells can

be removed for a composition simultaneously by using multiple antibodies with differing specificities. These antibodies may be localized in different layers of a multiple layer polymeric matrix. In this embodiment, the current applied to the different layers may be modulated so as to release each type of cell at a separate time thereby facilitating the efficient purification and separation of these cells.

The subject invention can also be used to isolate targets such as virus particles, proteins, polynucleotides, and the like, using molecules incorporated in the polymer matrix that bind to the target moiety. The targeted material is contacted with the polymer and then washed to remove unbound material. Bound material can be released according to the methods of the invention.

Although antibodies are exemplified herein as molecules capable of binding to a target moiety, other types of molecules can also be used with the subject invention. For example, cell adhesion molecules, lectins, ligands, receptors and the like can also be incorporated into a conductive polymer. In addition, polynucleotide molecules can be immobilized in a polymer matrix and used to selectively bind other polynucleotides, or even proteins, depending on the binding specificity and sequence of the polynucleotide.

Following are examples which illustrate procedures for practicing the invention. These examples should not be construed as limiting. All percentages are by weight and all solvent mixture proportions are by volume unless otherwise noted.

#### Example 1 – CD34<sup>+</sup> Cell Binding as a Function of Polymer Type and Antibody Concentration

It is difficult to obtain human hematopoietic stem cells. Therefore, ML-1 cells, a human non-adherent cell line expressing CD 34, was used to determine the capacity and specificity of the interaction between CD34<sup>+</sup> cells and immunopolymers.

ML-1 cells were grown in Iscove's modified Dulbecco's medium (IMDM) supplemented with 10% fetal bovine serum (FBS) and antibiotics (50 U/ml penicillin and 50

U/ml streptomycin). Cells were grown to confluence, then harvested by centrifugation, washed and resuspended at a concentration of  $10^5$  cells/ml.

The antibody used to give specificity to the conductive polymer was an anti-CD34 monoclonal antibody (HPCA-1, supplied at 50  $\mu$ g/ml in phosphate buffered saline with gelatin and 0.1% sodium azide, Becton Dickinson, CA). The antibody was stored at 4°C.

Two types of polymers were prepared. The polymers were prepared as follows:

Polymer 1: Polypyrrole doped with naphthalene sulfonate.

The following solutions were combined into a 50 ml beaker:

20 ml of a 0.08 M aqueous solution of 2-naphthalene sulfonate (pH 2.0)

1.5 ml of freshly distilled pyrrole (0.2 M final concentration)

various masses of anti CD34 antibody (0, 5, 10, and 20  $\mu$ g).

Polymer 2: Polypyrrole doped with p-toluene sulfonate.

The following solutions were combined into a 50 ml beaker:

20 ml of a 0.1 M p-toluene sulfonic acid dimethyl ammonium salt

1.5 ml of freshly distilled pyrrole (0.2 M final concentration)

various masses of anti CD34 antibody (0, 5, 10, and 20  $\mu$ g).

For each type of conductive polymer, reagents were added and mixed, then polymerized in a polymerization chamber with an indium tin oxide-coated glass anode and a platinum cathode, under a current of 1.6 V for 30 seconds followed by 1.0 V for 20 minutes. After polymerization, the polypyrrole membranes were removed from the anode, washed with IMDM and transferred into wells of 6-well culture plates containing 3 ml of complete medium each.

CD34<sup>+</sup> cells ( $10^5$  in 1 ml complete IMDM/well) were added to the polypyrrole membranes and incubated for 1 hour at 37°C in a non-gassed incubator. Following incubation, the membranes were removed from the cell suspensions, washed, then stained

using a modified Wright-Giemsa stain. The membranes were dried and fixed to a glass slide under a quick dry mounting medium and observed under light microscopy for the following parameters: number of bound CD34<sup>+</sup> cells, uniformity of binding, and membrane thickness.

In these experimental conditions, non specific binding of CD34<sup>+</sup> cells to the polymers was insignificant. When anti-CD34 antibody was present during the polymerization reaction, both types of conductive immunopolymers bound CD34<sup>+</sup> cells. However, for equal amounts of antibody present, the naphthalene-doped polypyrrole showed a higher binding capacity than the p-toluene-doped polymer (Figure 2). Thickness of the polypyrrole membranes influenced their binding capacity since more CD34<sup>+</sup> cells were bound in the thinner areas of membranes (Figure 3).

#### Example 2 – Effect of Polymerization Time on Cell Binding Capacity

The objective of this experiment was to determine the optimal polymerization time in the presence of antibody to obtain the highest cell binding capacity for the polypyrrole membranes.

ML-1 cells were grown in Iscove's modified Dulbecco's medium (IMDM) supplemented with 10% fetal bovine serum (FBS) and antibiotics (50 U/ml penicillin and 50 U/ml streptomycin). Cells were grown to confluence, then harvested by centrifugation, washed and resuspended at a concentration of 10<sup>5</sup> cells/ml.

The antibody used to give specificity to the conductive polymer was an anti-CD34 monoclonal antibody (HPCA-1, supplied at 50 µg/ml in phosphate buffered saline with gelatin and 0.1% sodium azide, Becton Dickinson, CA). The antibody was stored at 4°C.

Two types of polymers were prepared. The polymers were prepared as follows:

Polymer 1: Polypyrrole doped with naphthalene sulfonate.

The following solutions were combined into a 50 ml beaker:

20 ml of a 0.08 M aqueous solution of 2-naphthalene sulfonate (pH 2.0)

1.5 ml of freshly distilled pyrrole (0.2 M final concentration)

various masses of anti CD34 antibody (0, 5, 10, and 20 µg).

Polymer 2: Polypyrrole doped with p-toluene sulfonate.

The following solutions were combined into a 50 ml beaker:

20 ml of a 0.1 M p-toluene sulfonic acid dimethyl ammonium salt

1.5 ml of freshly distilled pyrrole (0.2 M final concentration)

various masses of anti CD34 antibody (0, 5, 10, and 20 µg).

For each type of conductive polymer, reagents were added and mixed, then polymerized in a polymerization chamber with an indium tin oxide-coated glass anode and a platinum cathode, under a current of 1.6 V for 30 seconds followed by 1.0 V for 20 minutes.

After polymerization, the polypyrrole membranes were removed from the anode, washed with IMDM and transferred into wells of 6-well culture plates containing 3 ml of complete medium each.

The cell binding capacity of the polymers increased with polymerization time. The higher cell binding capacity of the naphthalene-doped polypyrrole compared to the p-toluene-doped polypyrrole was already apparent after 5 minutes confirming the result obtained in the first experiment (Figure 4). The binding capacity of the naphthalene-doped polymer did not increase between 10 and 15 minutes suggesting that polymerization and antibody incorporation were complete in 10 minutes.

### Example 3 - Capacity and Specificity of Binding

NSC1.1 cells, a human non-adherent cell line expressing high levels of the CD 34 antigen, was used to determine the capacity and specificity of interaction between CD34+ cells and the polyphenol polymers. NSC1.1 cells were cultured in Iscove's modified Dulbecco's medium (IMDM) supplemental with 10% fetal bovine serum (FBS) and antibiotics (50 U/ml penicillin and 50 U/ml streptomycin); at 37°C in an humidified atmosphere of 5% CO<sub>2</sub> in air. Cells were harvested by centrifugation (4 min. at 300xg),

washed using complete growth medium and resuspended at a concentration of  $10^5$  cells/ml in Iscove's complete medium. To determine the specificity of cell binding, a B lymphocyte cell line characterized as  $CD34^+$  and  $CD19^+$  was used. This cell line was derived from a normal bone marrow aspirate. The B cell line was cultured under the same conditions as the NSC1.1 cells. The antibodies selected to test the binding specificity to the polyphenol conducting polymer was an anti- $CD34$  monoclonal antibody (HPCA-1) and a anti- $CD19$  antibody (Becton Dickinson Immunocytometry Systems, CA). Both these antibodies were supplied at 50  $\mu$ g/ml in phosphate buffered saline with gelatin and 0.1% sodium azide.

Studies using anti- $CD34$  antibodies entrapped in conducting polypyrrole electrode matrices have been successfully carried out. Table 2 demonstrates the binding of  $CD34^+$  cells to 2 different polypyrrole matrices containing anti- $CD34$  antibodies. These results indicate that, in the absence of antibodies in the system, the polypyrrole polymers do not bind  $CD34^+$  cells (therefore there is no non-specific binding). Membranes that have antibody (anti- $CD34$ ) incorporated into the polymer matrix do capture  $CD34^+$  cells (specific binding) and furthermore, the binding capacity of the immunopolymers increases proportionally when the concentration of anti- $CD34$  antibodies entrapped in the polymers is increased.

**Table 1.**  $CD34^+$  cell selection by electrochemically synthesized immunopolymers.

Antibody Concentration	Polymer Matrix/ $CD34^+$ Cells Bound <sup>1</sup>	
	Naph. Sulfonate Doped PolyPyrrole	p-Toluene Sulfonate Doped PolyPyrrole
0	0 ( $\pm 0$ ) <sup>2</sup>	0 ( $\pm 0$ )
5	82 ( $\pm 11$ )	48 ( $\pm 15$ )
10	556 ( $\pm 44$ )	116 ( $\pm 52$ )
20	814 ( $\pm 104$ )	210 ( $\pm 99$ )

<sup>1</sup>=approximately  $10^3$   $CD34^+$  cells were added to a 2 cm<sup>2</sup> polymer membrane

<sup>2</sup>=S.E. of the mean for 6 replicates



#### Example 4 – Specificity of Binding

To test the specificity of interaction between cells and conductive immunopolymers, antibodies with two different specificities (anti-CD19 and anti-CD34) were incorporated into separate membranes. These immunopolymers specific for CD34 or CD19 were tested for their ability to specifically bind to CD34<sup>+</sup> or CD19<sup>+</sup> cells.

The ML-1 cell line were used as a source CD34<sup>+</sup>, CD19<sup>+</sup> cells. Wilkinsons cells, a B-cell lymphoma clinical isolate from a bone marrow aspirate, were used as a source of CD34<sup>+</sup>, CD19<sup>+</sup> cells. CB-1 cells, a primitive neuro-ectodermal tumor line, were used as source of CD34<sup>+</sup>, CD19<sup>+</sup> cells. ML-1, Wilkinsons and CB-1 cells were grown in Iscove's modified Dulbecco's medium (IMDM) supplemented with 10% fetal bovine serum (FBS) and antibiotics (50 U/ml penicillin and 50 U/ml streptomycin). Cells were grown to confluence, then harvested by centrifugation, washed and resuspended at a concentration of 10<sup>5</sup> cells/ml.

The antibodies used were anti-CD34 monoclonal antibody (HPCA-1, supplied at 50 µg/ml in phosphate buffered saline with gelatin and 0.1% sodium azide, Becton Dickinson, CA), and anti-CD19 monoclonal antibody (Becton Dickinson, CA).

A naphthalene-doped polypyrrole was used to assess binding specificity. The following solutions were combined into a 50 ml beaker:

20 ml of a 0.08 M aqueous solution of 2-naphthalene sulfonate (pH 2.0)

1.5 ml of freshly distilled pyrrole (0.2 M final concentration)

various masses of anti CD34 antibody (0, 5, 10, and 20 µg).

Naphthalene and pyrrole were thoroughly mixed for 1 minute, then the magnetic stirrer was switched off and the polymerization was initiated using a current of 1.6 V for 30 seconds followed by 1.0 V for 10 minutes. Polypyrrole membranes containing anti-CD34 or anti-CD19 or no antibody were prepared. After polymerization, the newly formed membranes were removed from the chamber, washed with IMDM and transferred into wells of 6-well culture plates containing 3 ml of complete medium each.

After incubation with ML-1 or Wilkinsons, or CB-1 cells, the number of membrane-bound cells were determined by light microscopy. Results are summarized in Table 2.

**Table 2.** Cell binding specificity of a naphthalene-doped conductive polymer. Data represent the number ( $\pm$  SE) of cells bound to naphthalene-doped polymers containing different types of antibodies (anti-CD34 or anti-CD19 or none).

Cell Type	Naphthalene-doped polypyrrole specificity		
	anti-CD34	anti-CD19	none
CD34 <sup>+</sup> /CD19 <sup>-</sup> (ML-1)	146 $\pm$ 29 <sup>1</sup>	1.5 $\pm$ 2.	(-) <sup>2</sup>
CD34 <sup>-</sup> /CD19 <sup>+</sup> (Wilk.)	(-)	8.6 $\pm$ 1.5	(-)
CD34 <sup>-</sup> /CD19 <sup>-</sup> (CB-1)	(-)	(-)	(-)

1= S.E. for the mean of 4 membranes (500 cells added/membrane + 10 $\mu$ g total antibody)

2= (-) no visible cells binding

Only cells expressing the CD marker recognized by each antibody type were able to bind the conductive immunopolymers suggesting that the interaction between cells and polymer is specifically mediated through the monoclonal antibodies included in the polymer.

#### Example 5 – Specific Selection of Rare CD34<sup>+</sup> Cells From a Large Excess of CD34<sup>-</sup> Cells

The objective of this study was to determine the ability of a naphthalene-doped immunopolymer to specifically bind rare CD34<sup>+</sup> cells in the context of a large excess of CD34<sup>-</sup> cells.

The ML-1 cell line was used as a source CD34<sup>+</sup>, CD19<sup>-</sup> cells. CB-1 cells, a primitive neuro-ectodermal tumor line, were used as source of CD34<sup>-</sup>, CD19<sup>-</sup> cells. ML-1 and CB-1 cells were grown in Iscove's modified Dulbecco's medium (IMDM) supplemented with 10% fetal bovine serum (FBS) and antibiotics (50 U/ml penicillin and 50 U/ml streptomycin). Cells

were grown to confluence, then harvested by centrifugation, washed and resuspended at a concentration of  $10^5$  cells/ml.

In order to visualize the two different cell populations, the  $CD34^+$  cells were stained dark blue using a Giemsa stain. The  $CD34^-$  cells (CB-1) were stained pale pink. Stained cells were enumerated and  $CD34^+$  cells (ML-1) were seeded into CB-1 cells so that ML-1 cells represent only 10% of all cells in the mixture.

The antibodies used were anti- $CD34$  monoclonal antibody (HPCA-1, supplied at 50  $\mu\text{g/ml}$  in phosphate buffered saline with gelatin and 0.1% sodium azide, Becton Dickinson, CA).

A naphthalene-doped polypyrrole was used to assess binding specificity. The following solutions were combined into a 50 ml beaker:

20 ml of a 0.08 M aqueous solution of 2-naphthalene sulfonate (pH 2.0)

1.5 ml of freshly distilled pyrrole (0.2 M final concentration)

Various masses of anti  $CD34$  antibody (0, 5, 10, and 20  $\mu\text{g}$ ).

Naphthalene and pyrrole were thoroughly mixed for 1 minute, then the antibody was added, and the mixture was stirred for an additional minute. Then, the magnetic stirrer was switched off and the polymerization was initiated using a current of 1.6 V for 30 seconds followed by 1.0 V for 10 minutes. After polymerization, the newly formed membranes were removed from the chamber, washed with IMDM and transferred into wells of 6-well culture plates containing 3 ml of complete medium each.

The cell mixture (90% CB-1 + 10% ML-1) was applied to the polymerized membranes and incubated for 30 minutes at  $37^\circ\text{C}$  in a non-gassed incubator. Following incubation, the membranes were removed from the cell suspensions, washed 3 times, air dried and fixed to a glass slide under a quick dry mounting medium. The slide-mounted membranes were observed under light microscopy to determine the color (dark blue versus pale pink) and the number of bound cells.

Based on color, the CD34<sup>+</sup> cells represented 7.6% of all cells. After 30 minutes of incubation, only dark blue cells (CD34<sup>+</sup>) were observed at the surface of the membranes. These results indicate that the naphthalene-doped polypyrrole can be used to select CD34<sup>+</sup> cells in a cell mixture containing a large excess of irrelevant cells.

#### Example 6 - Recovery and Viability of Target Cells

The ability of the subject invention to capture and release a target cell population was demonstrated using a purified CD34<sup>+</sup> cell population as the starting material. The polymer was prepared with anti-CD34 antibody incorporated therein as described previously.

**Table 3.** Recovery and viability of CD34<sup>+</sup> selected cells.

	Mean Cell Count (CD34 <sup>+</sup> )	Mean Percent
# of CD34 <sup>+</sup> cells added	500	
CD34 <sup>+</sup> Cells Bound	448 ± 14 <sup>1</sup>	90 ± 2
CD34 <sup>+</sup> Cells Released/Recovered	378 ± 12	84 ± 1
Viability of CD34 <sup>+</sup> cells Recovered	362 ± 6	96 ± 2

(1): S.E. of the mean for 4 membranes

(2): 10µg total antibody was added to the polymer mixture

The ability to specifically capture rare cells, such as HSC, from a complex mixture of irrelevant cells (peripheral blood, and/or bone marrow) is an important requirement for efficient cell selection. Non-specific binding is a critical factor limiting the specificity of the current marketed devices. The conducting properties of this type of polymer can be exploited to minimize this phenomenon by modulating the net charge of the immunoaffinity surface in order to prevent non-specific binding while allowing specific antibody-based cell binding. Therefore, the ability of a conductive polymer of the subject invention to be able to capture targeted cells from a complex tissue such as peripheral blood or bone marrow was examined. CD34<sup>+</sup> cells were seeded at varying concentrations (10%, 5% and 1%) into freshly isolated

peripheral blood leukocytes (PBL) and the mixture applied to an immunopolymer comprising anti-CD34 antibody (10 $\mu$ g total anti-CD34 antibody). The results from one of those experiments is shown in Table 4. The data shows that a conductive polymer of the present invention is capable of selectively binding to a target cell population (in this study CD34<sup>+</sup>) and recovering a high percentage of cells with an extremely high degree of purity (see Table 4). These data strongly suggest that the present invention is capable of performing to a high level of accuracy (specific target cell recovery) when applied to a complex cell suspension and can selectively capture stem cells at concentrations equivalent to those found in clinical relevant samples of mobilized peripheral blood (1-5%).

**Table 4.** Selective binding and percent capture of CD34<sup>+</sup> cells from peripheral blood leukocytes.

% CD34 <sup>+</sup> Cells Recovered From PBL (Mean of 6 membranes)			
% CD34 <sup>+</sup> Cells/PBL	CD34 <sup>+</sup> Recovered	Purity	Viability
10%	83%	89.2%	96%
5%	93%	93.5%	95%
1%	70%	94.4%	93%

A donor bone marrow harvest ranges on average between 700-1000 ml resulting in a total nucleated cell count of approximately 1-2x10<sup>10</sup>. Assuming an average of 1% CD34<sup>+</sup> in normal marrow, the total CD34<sup>+</sup> count would average 1-2x10<sup>8</sup>. It is accepted that a minimum of 1x10<sup>6</sup> cell/Kg is required for CD34<sup>+</sup> cell transplantation. Therefore, a transplant patient with an average weight of 75Kg required 75x10<sup>6</sup> CD34<sup>+</sup> cells for a successful transplant. This is compared to 10<sup>8</sup> Kg total nucleated bone marrow cells that would be required for a similar bone marrow transplant. Thus, the purity (93%-95%) and cell recovery rate (70%-93%) obtained using a conductive immunopolymer according to the present invention is adequate to give the required concentration of cells for transplantation.

### Example 7 - Indirect Entrapment of Anti-CD34 Antibody in the Polymer via Immobilized NeutrAvidin

NEUTRAVIDIN is immobilized in a polyphenol matrix under optimal conditions of accessibility. NEUTRAVIDIN (Pierce, Rockford, IL) is a biotin binding protein that lacks both the carbohydrate domain of avidin and the RYD (fibronectin mimic) cell surface receptor binding domain of streptavidin. Both of these domains can lead to significant non-specific cell binding levels.

Either the biotin-anti-CD34 antibody complex can be bound to the immobilized, surface-exposed NEUTRAVIDIN, followed by CD34<sup>+</sup> cell binding; or the CD34<sup>+</sup> cell population can be pre-mixed with and bound to biotin-anti-CD34 antibody, followed by binding of the biotin-anti-CD34 antibody-CD34<sup>+</sup> cell complex to surface immobilized and accessible NEUTRAVIDIN in the polyphenol matrix. With either of these two approaches, a number of biotin derivatives of anti-CD34 antibody can be used. These derivatives are described below. Their advantages are: (1) that they provide relatively mild cleavage chemistries, making possible anti-CD34 antibody; CD34<sup>+</sup> cell release; (2) they are long chain derivatives allowing for closer contact between the polyphenol matrix and the bound CD34<sup>+</sup> cell; and (3) they allow for better orientation of the anti-CD34 antibody (antigen-binding region oriented outward) in the selection system matrix.

### S-S- Cleavable Biotins

NHS-SS-Biotin and Biotin-HPDP are two reagents (Pierce, Rockford, IL) containing -S-S- bonds. Both are long chain length reagents (24.3 and 29.2 Å respectively). NHS-SS-Biotin will react with amine groups on the antibody. This biotinylated anti-CD34 antibody species interacts with NEUTRAVIDIN in the normal way, but the -S-S- bond can be chemically cleaved under reducing conditions (50 mM DTT or 100 mM β-mercaptoethanol) releasing the antibody bound to the CD34<sup>+</sup> cell. Biotin-HPDP is a unique reagent that can react with sulfhydryls. If the CD34 antibody is first reduced under mild conditions with

mercaptoethylamine-CDI in EDTA, the antibody will be cleaved at the level of disulfide bonds between the two heavy chains generating -SH groups on each heavy chain. This mild reduction treatment will retain the antigen-specific binding capacity of the antibody. At this point, the Biotin-HPDP may be reacted against these -SH groups in the hinge region of the molecule. The reaction is complete within 90 minutes in a range of buffers pH 6-9. Following binding of the antibody to CD34<sup>+</sup> cells, this complex will be added to the polyphenol matrix. Once bound to the polymer, the biotin can be cleaved and removed with either 100 mM  $\beta$ -mercaptoethanol or 50 mM DTT (Pierce, Rockford, IL) to release the cells.

#### NHS Iminobiotin and Iminobiotin Hydrazide

These derivatives of biotin (the guanidino analog) have the interesting and useful properties that at neutral to higher pH (maximum binding level above pH 9) they bind normally to the NEUTRAVIDIN binding site. When the pH is lowered below pH 6, their affinity is gradually reduced, and at pH units close to 4, it is totally dissociated. Alternately, the complex can be completely dissociated at any pH by competition with 1 mM biotin for 15 min. (Zeheb *et al.*, 1983). Therefore, in these studies, the anti-CD34 antibody will be derivatized either at a primary amine with NHS Iminobiotin under the conditions described by the supplier (Pierce, Rockford, IL) or at a sialic acid residue in the carbohydrate moiety reacted first with 1 mM NaIO<sub>4</sub> (at 0°C) to form the aldehyde, followed by reaction with Iminobiotin hydrazide (BioAffinity Systems, Inc.) To form the hydrazone stable from pH 2-10. This derivative should result in an anti-CD34 antibody with its antigen binding site oriented outward from the matrix immobilized NEUTRAVIDIN, thus increasing the probability of CD34<sup>+</sup> cells binding the anti-CD34 antibody. Conversely, when the derivatized antibody is bound to CD34<sup>+</sup> cells first, there will be a greater probability of the Iminobiotin moiety, (oriented near the Fc portion of the antibody molecule) binding preferentially to the polyphenol matrix immobilized, NEUTRAVIDIN. Biotin LC-Hydrazide (Pierce, Rockford, IL), a commercially available long chain (24.7 Å) biotin reagent can be used as a substitute

for the biotin derivative mentioned above. In all cases, antibodies have been shown to retain full immunological activity following this biotinylation, since the derivatization of carbohydrate resides in the Fc portion of the antibody molecule, distal to the antigen binding site. To obtain strong binding of the Iminobiotin moiety to immobilized NEUTRAVIDIN, the pH will be maintained at pH 8.5 or above.

#### Example 8 - Monitoring Systems to Assess Maximum Cell Binding/Release, and the Physiologic State of Bound Cells

Exemplified herein are two classes of electrochemical measurements which can be performed to monitor cell binding, release and viability. The first method involves monitoring of change in electrochemical impedance or the related electrochemical parameters (Resistance, capacitance). When cells bind to the conducting immunopolymer via the antibodies, the dielectric constant thickness of the dielectric layer, can change in the vicinity of the electrode surface. At a constant frequency (usually 1-10 KHz), a gradual increase of the impedance could be detected after addition and binding of the cells. The magnitude of the change is typically proportional to the number of bound cells. Kinetics of the cell binding will be followed by continuous monitoring of the impedance. Small platinum or gold interdigitated "comb" electrodes can be used. Impedance can be monitored using a potentiostat coupled with a lock-in amplifier controlled with a computer through an IEEE-488 interface driven by electrochemical impedance software. Preferably, the polymer film is kept very thin ( $\leq 100\text{\AA}$ ) to ensure adequate sensitivity to detect cell binding.

A second monitoring system is the enzyme-linked amperometric immunoassays. The microcurrent produced by entrapped reporter enzymes (HRP and glucose oxidase-(GOx)) will sense glucose. This sensing is carried out experimentally by coupling the enzymes HRP and GOx as shown in Figure 5.

Glucose is reflected as a microcurrent,  $1(\mu\text{A})$ , necessary to regenerate the HRP enzyme following cycling through the two-step reaction sequence (Figure 5). For calibration



experiments, different concentrations of glucose can be added to the solution without cells, the current reflecting the production of  $H_2O_2$  at a controlled potential - 0.05 V wrt Ag/AgCl (previously shown to be a safe potential for cells), measured to establish a calibration curve. The current will change as the cells bind due to either the local consumption of glucose or the prevention of it diffusing into the enzyme immobilized electrode by the bound cell. Thus, the change in current will also be capable of calibration and conversion into the number of bound cells. Dynamic curves of current Vs time can be investigated as a function of the adsorption/desorption of  $CD34^+$  stem cells seeded at varying concentrations into PBL.

Not only does the latter monitoring system allow for the detection of bound cells and their subsequent release but their viability is also assessable through the glucose measurements. Viable cells possess basal metabolic rates which are capable of calibration by this system. Also, by utilizing other oxidase enzymes, in combination with HRP, other substrate may be assayed that also reflect cell viability.

It should be understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and the scope of the appended claims.

References

Alva, K.S., K.A. Marx, L.A. Samuelson (1996) *S.P.I.E.*, 2716:152.

5

Bender *et al.* (1991) *Blood*, June, 77:2591-2596.

Berenson, R.J., W.J. Bensinger, R.S. Hill (1991) *Blood*, 77:1717.

de Wynter, E.A., L.H. Coutinho, X. Pei (1995) *Stem Cells*, 13:524.

10

Zeheb, R., V. Chang, and Orr Ga (1983) *Anal. Biochem.*, 129:156.

U.S. Patent No. 4,714,680.

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
215